



# New York

## Live births, abortions, and pregnancies


Live birth rates, 1982–1992



Live birth rates,\* 1991



Pregnancy rates, 1990



National live birth rate ranking, 1990


Age group		
15-17	18-19	15-19
36th highest	43rd highest	40th highest

\*This graph refers to females aged 15–19 years.




# Sexually transmitted diseases New York

## Gonorrhea rates,\* 1982–1992



## Gonorrhea rates, 1992



## National gonorrhea rate ranking,<sup>†</sup> 1992

Males – 26th highest    Females – 24th highest


\*Data were unavailable before 1985.

†National ranking refers to males and females aged 10–24 years.




## Chlamydia cases and rates,\* 1988–1992


Cases




Rates





Cases



Rates



## Acquired immunodeficiency syndrome (AIDS) cumulative cases, 1981–1992



\*Chlamydia cases and rates refer to males and females of all ages. Data for males were unavailable before 1989.



# Related risk behaviors,\* 1993

New York

Had sexual intercourse (lifetime)

*Question not included in survey.*

Had sexual intercourse with four or more people (lifetime)

*Question not included in survey.*

Had sexual intercourse during the past three months (currently sexually active)

*Question not included in survey.*


Used a condom during last sexual intercourse (among currently sexually active)

*Question not included in survey.*

Used birth control pills during last sexual intercourse (among currently sexually active)

*Question not included in survey.*

Injected illegal drugs (lifetime)



\*Related risk behaviors refer to high school students.

Sources: CDC—NCHS and NCCDPHP (birth, abortion, and pregnancy data), NCPS (gonorrhea and chlamydia data), NCID (AIDS data); 1993 New York Youth Risk Behavior Survey (risk behavior data).

